

Botanical Ratchets

I.M. Kulić¹, M. Mani¹, H. Mohrbach², R. Thaokar³ and L. Mahadevan^{1,4}

¹ *Engineering and Applied Sciences, Harvard University, Cambridge, MA 02138, USA*

² *Physique Moléculaire et des Collisions, Université Paul Verlaine-57012 Metz, France,*

³ *Chemical Engineering, IIT Bombay, Mumbai 400076 India, and*

⁴ *Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA*

Ratcheting surfaces are a common motif in nature and appear in plant awns and grasses. They are known to confer selective advantages for seed dispersion and burial. In two simple model experiments we show that these anisotropically toothed surfaces naturally serve as motion rectifiers and generically move in a unidirectional manner when subjected to temporally and spatially symmetric excitations of various origins. Using a combination of theory and experiment we show that a linear relation between awn length and ratchet efficiency holds under biologically relevant conditions. Thus, grass awns efficiently transform non-equilibrium environmental stresses into useful work and directed motion using their length as a fluctuation amplifier, yielding a selective advantage to these organelles in many plant species.

PACS numbers: 87.18.Tt 87.18.Nq 87.10.-e 87.19.ru

Motion rectification by ratchets arises repeatedly as a theme with variations in nature to technology on a range of scales, from the molecular [1]-[6] to the macroscopic [7]-[9] in both deterministic and stochastic systems. A key ingredient in all ratchets is the presence of a static or dynamic asymmetry that serves to rectify motion, with many convergent solutions seen in nature. For example, ratchet and pawl mechanisms are basic mechanical components in many man-made gadgets and machines. They serve to rectify linear or rotational motions by using a mechanical anisotropy that often takes the form of an asymmetric saw-tooth that allows the ratchet to slip in one direction and not in the opposite direction. Their microscopic (thermal) analogs have been intensely studied [1]-[6] over the past decades in context of molecular motors [5], where the physical sawtooth is replaced by a potential with a similar property that is periodically switched on and off. On the macroscopic scale as well, nature has stumbled on ratchets repeatedly, and a host of examples exist in the plant world in particular [10]-[14], where they might confer selective advantages for seed dispersion and burial. Indeed, the fact that different plant families show the same robust and efficient solution for the dispersion and self burial problem suggests that this is yet another example where a simple physical principle often leads to convergent evolution. Here we use the particular example of the wild foxtail *Hordeum murinum* to study macroscopic natural ratchets that rectify deformation induced by swelling, shrinking, and other environmental factors into unidirectional motion that serves important functional purposes, from the burial of seeds to their transport by animals.

When a *Hordeum murinum* plant awn is placed on a soft, porous or rough material and excited by macroscopic forces with a variety of origins, it generically moves in the direction of its tip. To understand the mechanism of operation of this simple machine, we start with the




FIG. 1: (Color online) a) Common grass species exhibiting ratchet motility (from left to right): single spikelet of foxtail grass (*Hordeum murinum*), *Setaria viridis* and barley (*Hordeum vulgare*), scale bar 2 cm. b) *Hordeum murinum*'s micro-barb surface structure responsible for ratchet effect (SEM), scale bar 50 μm . c) A single clamped unstrained *Hordeum murinum* awn (with micro-barbs pointing to the right) glides easily on substrate moving to the right (paper cover). d) Awn buckling and progressive micro-barb locking induced by substrate motion to the left. e) A barb-locked awn on substrate resists also upward lift forces.

common observation that sliding one's fingertips over the surface of a foxtail awn is easy in one direction and essentially impossible in the opposite direction. A closer look under a scanning electron microscope (SEM) reveals the reason (Fig 1b). The awn has a highly anisotropic surface structure with sharp micro-barbs tilted unidirectionally at an angle $\beta \approx 35^\circ$ with the horizontal and a typical length of 50 μm . Such a surface geometry in combination with the elasticity of the awn induces a cooperative locking transition of micro-barbs via a progressive bending of the awn toward the substrate (Fig 1d). This cooper-




FIG. 2: (Color online) a) Foxtail grass moves unidirectionally on an harmonically oscillating horizontal shaker (cf. supplementary movie 1). b) The mean velocity is shown as a function of the shaker frequency, with error-bars from 5 different foxtail specimen (circles). The best fit to a theoretical model (see text) with $\gamma = 0.4$ is shown as a thick line. Inset: Graphical model for ratchet motion on oscillating substrate. The ratchet leaves the substrate frame when its inertial force overcomes the friction force at $t = t_1$ and it locks and moves with the substrate again when ratchet and substrate velocities coincide again at $t = t_2$.

activity of locking increases the number of stress bearing micro-barbs and thus reduces the probability of interfacial failure. The tilted barb has a second effect in addition to providing a force parallel to the substrate, since its geometry naturally also gives rise to an effective normal (adhesion-like) force component $F_{\perp} = F_{\parallel} \tan \beta$ as shown in Fig 1e. The combination of a rigid microscopic geometry associated with the barbs and a macroscopic elastic compliance associated with the backbone to which the barbs are attached allows the foxtail awn to slide unhindered with respect to the substrate in one direction (Fig 1c) but efficiently suppresses any motion in the opposite direction (Fig 1d). We consider now two different physical situations in which transport is generically observed.

Transport in inertially dominated environments. In a first experiment we placed *Hordeum murinum* spikelets on a horizontal shaker driven sinusoidally with amplitude $A = 3.7\text{mm}$ and observe their motion as a function of the driving frequency ω . There is a critical frequency

ω_c below which almost no motion is observed. Above the critical shaking frequency we observe the onset of strong spikelet drift in the direction of its tip, as shown in Fig 2 (See supplementary movie 1 [17]). The foxtail cannot move unless inertial forces overcome friction, so that we expect the onset of motion when $\omega > \omega_c$ with $\omega_c^2 mA = \gamma mg$, i.e. $\omega_c = \sqrt{\gamma g/A}$. Here m is the mass of the grass, γ its friction coefficient and g the acceleration due to gravity. Above this critical frequency, the mean transport velocity can be derived from the following graphical argument (cf. inset of Fig 2b): the grass moves with the shaker up to the point t_1 where the shaker acceleration $\dot{v}_s(t)$ coincides with the critical slope $-\gamma g$. From this point on the grass unbinds from the shaker and slides on it, decelerating under the influence of friction with $\dot{v}(t) = -\gamma g$. The free motion ceases when the grass and substrate velocities coincide again at a later time $t = t_2$ and the cycle begins again. In general the time-averaged grass velocity can be written as $\bar{v} = \frac{1}{T} \int_{t_1}^{t_2} (v_s - \gamma g t) dt$. Evaluating this expression for a sinusoidal shaker velocity $v_s(t) = A\omega \sin \omega t$ shows that this agrees well with our experimental results, with $\gamma = 0.4$ (Fig 2b), which value is close to the experimentally measured friction coefficient of $\gamma = 0.46 \pm 0.05$ (from independent inclined plane measurements on 6 different *Hordeum murinum* samples). Close to the transition frequency ω_c , a Taylor expansion for t_1 and t_2 in the small parameter $\alpha = \sqrt{1 - \omega_c^2/\omega^2} < 1$ gives simple expressions $t_1 \approx \frac{T}{2} - \sqrt{2}\alpha/\omega_c$ and $t_2 \approx \frac{T}{2} + 2\sqrt{2}\alpha/\omega_c$, leading to $\bar{v} = \frac{1}{T} \int_{t_1}^{t_2} (v_s - \gamma g t) dt \approx \frac{9}{2} \Theta(\omega - \omega_c) A\omega \left(1 - (\omega_c/\omega)^2\right)^2$ with Θ the Heaviside function. In the other limit of large frequencies $\omega \gg \omega_c$ the grass moves almost freely obtaining momentum from the shaker during a very brief locking phase, so that we expect an asymptotic mean velocity $\bar{v} \approx A\omega$. Summarizing, the transport velocity in this frictional-inertial regime is therefore independent of grass mass and length but obeys a non-linear behavior at onset that transitions to a simple linear relation far from criticality.

Transport in continuously deforming environments. We now turn to the second, biologically more relevant, situation in which net transport is observed: continuous deformation of the substrate along which the grass is moving. Two important examples for such deformations are: a) Deformation of the skins of animals (naturally dispersing the foxtail) b) Diurnal and seasonal soil swelling and shrinkage driven by humidity and temperature variations [15],[16]. The first process is believed to be responsible for the dispersal of foxtail over large distances. It poses a hazard to animals if the foxtail is trapped in nostrils or ear channels, eventually ratcheting through the tissue causing infection and possibly death. The second process is a potential mechanism for the seed and awn burial into the soil [10]-[14], an important precursor of germination. In general in addition to substrate deformation the contour length of the ratchet grass itself

can actively vary as well. For example, relatives of the foxtails like *Stipa somata* show a hygroscopically driven coiling and kinking of their awns to vary the effective contour length of awns (along the direction of motion) which yields linear strains of up to 20% [10]. In either case, these strain variations may be rectified in the direction of the grass tip. To understand the minimal properties of ratchet propulsion in this case, we consider the combination of a time-periodic hygroscopic strain of the grass awn $\varepsilon_g(t)$, and that of the surrounding substrate (soil/animal fur) $\varepsilon_s(t)$. Then, on dimensional grounds, we expect a simple scaling relation of the mean propulsion velocity of the awn:

$$\bar{V}_g \sim aT^{-1}f(\varepsilon_s, \varepsilon_g) \quad (1)$$

where a is the awn length, T the period of the soil and awn strain variation and $f(\varepsilon_s, \varepsilon_g)$ a dimensionless functional.

For a more careful analysis we parameterize the grass using its unstrained arclength $s_g \in [-a/2, a/2]$ with $s_g = 0$ corresponding to its center. In response to internal and substrate strain, each point along the grass moves so that its position is $x_g(s_g, t) = X_g(t) + s_g(1 + \varepsilon_g(t))$ and its velocity is $v_g(s_g, t) = \dot{X}_g(t) + s_g\dot{\varepsilon}_g(t)$ with $X_g(t) = x_g(0, t)$ and $v_g(s, t) = \dot{x}_g(s, t)$. The substrate (soil tube) can be parameterized in a similar manner via its unstrained arclength coordinate s_s so that material points along it have a position $x_s(s_s, t) = (1 + \varepsilon_s(t))s_s$ and velocity $v_s(s_s, t) = s_s\dot{\varepsilon}_s(t)$. At a given instant of time t the grass is in contact with the soil over the interval $x_s \in [X_g(t) - \frac{a}{2}(1 + \varepsilon_g(t)), X_g(t) + \frac{a}{2}(1 + \varepsilon_g(t))]$. Then, the relative velocity between the grass and the substrate parameterized in terms of the grass material parameter s_g is $V_{rel}(s_g, t) = v_g(s_g, t) - v_s(s_s(s_g, t), t)$. We assume that the grass acts as an ideal ratchet, i.e. that no part of it can move in the *negative* direction (direction of the micro-barb sharp tip orientation) with respect to the contact points with the substrate. Then the relative velocity has to satisfy the following physical condition:

$$\min_{s_g \in [-\frac{a}{2}, \frac{a}{2}]} V_{rel}(s_g, t) = 0 \quad (2)$$

Using the fact that $x_s(s_s, t) = x_g(s_g, t)$ and $\min_{s_g \in [-a/2, a/2]} (s_g C) = -\frac{a}{2}|C|$ (for any value C) the upper condition can be recast as an equation of motion

$$\frac{d}{dt} \frac{X_g}{1 + \varepsilon_s} = \frac{a}{2} \frac{|(1 + \varepsilon_s)\dot{\varepsilon}_g - (1 + \varepsilon_g)\dot{\varepsilon}_s|}{(1 + \varepsilon_s)^2} \quad (3)$$

Integrating the previous expression yields the position of the center of the awn $X_g(t) = \frac{a}{2}(1 + \varepsilon_s) \int_0^t (1 + \varepsilon_s)^{-2} |(1 + \varepsilon_s)\dot{\varepsilon}_g - (1 + \varepsilon_g)\dot{\varepsilon}_s| dt'$. To determine the mean velocity of the awn, we assume that both $\varepsilon_s(t)$ and $\varepsilon_g(t)$ are simple periodic functions with a single minimum and maximum, with same




FIG. 3: (Color online) Ratchet grass propulsion in a continuously deforming environment. a)-b) Experimental setup: foxtail spikelets are placed inside a 4 mm inner diameter rubber tube that is periodically strained (see supplementary movie 2). Spikelets move in the direction of their tips (stained black for video analysis) with efficiency varying with their length. c) An intensity line-scan (kymograph) corresponding to the time dynamics in a) showing the periodic tube motion and the average propulsion of the spikelet over the time course of 60 seconds (x-axis). d) Propulsion distance per strain cycle vs. maximal rubber tube strain for 4 spikelets of various lengths: Experimental points with error bars from 7-9 measurements and best fits to $\Delta_g^I = a\varepsilon_s^{\max}/(1 + \varepsilon_s^{\max})$ (thick lines). Inset: The fitted effective contact length (y axis, units of cm) vs. the measured length of the longest awn (x axis, units of cm) of individual spikelets. While the effective lengths show a good correlation with the measured lengths, they are slightly smaller due to occasional awn buckling and imperfect tube-awn contact.

period T and that ε_s satisfies $\varepsilon_s(0) = \varepsilon_s(T) = 0$. Then the velocity averaged over a complete cycle $\bar{V}_g = (X_g(T) - X_g(0))/T$ is

$$\bar{V}_g = \frac{a}{2T} \int_0^T \frac{|(1 + \varepsilon_s)\dot{\varepsilon}_g - (1 + \varepsilon_g)\dot{\varepsilon}_s|}{(1 + \varepsilon_s)^2} dt \quad (4)$$

There are two interesting limiting cases worth considering: *Case I: Negligible substrate strain.* In this case $\varepsilon_s = 0$ so that (4) simplifies to $\bar{V}_g = \frac{a}{2T} \int_0^T \dot{\varepsilon}_g dt = \frac{a}{2T} (\int_0^{T_{\max}} \dot{\varepsilon}_g dt - \int_{T_{\max}}^T \dot{\varepsilon}_g dt) = \frac{a}{T} \varepsilon_{\max}$, where $T_{\max}$ is the point at which ε_g attains its maximum and $\dot{\varepsilon}_g$ changes sign from positive to negative. Then the propulsion distance per cycle

$$\Delta_g^I = T\bar{V}_g = a\varepsilon_{\max} \quad (5)$$

with $\varepsilon_{\max}$ being the maximal strain during a single strain cycle. *Case II: Negligible plant awn strain.* In this case we have $\varepsilon_g = 0$ and $\bar{V}_g = \frac{a}{2T} \int_0^T [|\dot{\varepsilon}_s| / (1 + \varepsilon_s)^2] dt$. Splitting the integral as in the previous case, we can perform the integration over a full cycle and obtain $\bar{V}_g = (a/T) \varepsilon_{\max} (1 + \varepsilon_{\max})^{-1}$ and

$$\Delta_g^{II} = a \frac{\varepsilon_{\max}}{1 + \varepsilon_{\max}} \quad (6)$$

We see that the displacement per cycle Δ_g scales in both cases linearly with the grass length a showing that longer grasses pick up and rectify larger fluctuations in proportion with their length. Therefore longer grass awns act effectively as better "fluctuation amplifiers".

To test our results quantitatively, we focus on the case when substrate strains dominate, since this is simple to setup experimentally. To simulate the behavior of a ratcheting awn in contact with soil, awns and single rachis spikelets of various grasses of different lengths and diameters (Fig 1 a) were placed inside a rubber tube (inner diameter 4 mm). The tube was then subjected to slow periodic strains with maximal strains ranging from $\varepsilon_{\max} = 0.01$ to 0.4. All the grasses rectified the oscillatory strain with the *Hordeum murinum* spikelets showing most robust and reproducible behavior (Fig 3 d). At very small sub-Hertz frequencies at which the rubber tubes were strained (0.3 Hz) the ratchet displacements per cycle were independent of both frequency and the direction of gravity. The experimental results for the propulsion displacement Δ_g as a function of the length of *Hordeum murinum* spikelets show good agreement with the theoretical prediction 6 with a single fitting parameter per specimen - the effective contact length a , which itself is proportional to the actual awn length (Fig 3 d, inset). For longer awns the fitted length shows however a systematic 20-40% reduction relative to the measured length consistent with our observation of occasional awn buckling and imperfect tube - awn contact of longer spikelets; this effect is less pronounced for shorter spikelets (Fig 3 d, inset).

Interestingly the results (5)-(6) are consistent with the previously measured linear relation between the awn length and burial depth of *Stipa somata* [13] and provide some evidence for a selective advantage of larger awn lengths. More generally, for hygroscopically driven awns (4) predicts a non-trivial interaction between soil strain and awn strain contributions to the rectification of motion. Indeed the two effects act in concert or "resonate"

for oppositely phased strains but can nullify each other for in-phase strains.

Our simple experiments and theory allow us to understand the physics of plant awn ratchets. Equipped with anisotropically toothed microstructured elastic awns the seeds are able to lock directionally into virtually any soft natural material, like soil, animal fur and skin. The pronounced asymmetry of their surface structures enables the seeds to rectify external environment fluctuations along the direction imposed by the specific orientation of their micro-barbs. Furthermore, longer awns proffer a simple selective advantage to those seeds since they can move further and thus disperse more efficiently than their shorter competitors. Our results also point the way for technological designs that might mimic this simple but robust motif.

-
- [1] Feynman Lectures on Physics, Addison-Wesley, Vol. 1, 1964.
 - [2] Ajdari A and J.R. Prost, C. R. Acad. Sci. II (Paris), 315, 1635 (1992)
 - [3] M.O. Magnasco, Phys. Rev. Lett. 71, 1477 (1993)
 - [4] P. Reimann, Phys. Rep. 361, 57 (2002)
 - [5] F. Julicher, A. Ajdari A, J.R. Prost, Rev. Mod. Phys. 69, 1269 (1997)
 - [6] I.M. Kulić, R.M. Thaokar and H. Schiessel, Europhys. Lett. 72, 527 (2005)
 - [7] A. Bug and B. Berne, Phys. Rev. Lett. 59, 948 (1987)
 - [8] B. Nordén, Y. Zolotaryuk, P. L. Christiansen, and A. V. Zolotaryuk, Phys. Rev. E 65, 011110 (2001)
 - [9] L. Mahadevan, S. Daniels and M. Chaudhury, Proc. Natl. Acad. Sci. (USA), 101, 23 (2003).
 - [10] L. Murbach, Botanical Gazette, 30, 113 (1900)
 - [11] M.H. Peart, J. Ecology, 67, 843 (1979)
 - [12] N.A. Stamp, J. Ecology, 72, 611 (1984)
 - [13] L.K. M. Garnier and I. Dajoz, Ecology, 82, 1720 (2001)
 - [14] R. Elbaum, L. Zaltzman, I. Burgert and P. Fratzl, Science 316, 884 (2007)
 - [15] S. Iwata, T. Tabuchi, B. P. Warkentin, Soil-Water Interactions, M. Dekker New York and Basel (1988).
 - [16] Soil Physics, T. J. Marshall, J. W. Holmes, and C. W. Rose, Cambridge University Press (1996).
 - [17] Online supplementary movie 1: *Hordeum murinum* moving on a laboratory shaker above the critical frequency. Recorded at 200 fps (frames/s), reproduced at 25fps.
 - [18] Online supplementary movie 2: *Hordeum murinum* spikelet moving in a stretching rubber tube. Recorded at 5 fps and reproduced at 25 fps. Color mark distance on tube in relaxed state = 2.54 cm.